

Forum

ON THE DETECTION OF LIPID HYDROPEROXIDES IN BIOLOGICAL SAMPLES

WILLIAM A. PRYOR

Biodynamics Institute, Louisiana State University, Baton Rouge, LA 70803-1800, U.S.A.

In 1987 this Journal published a Forum on the detection of lipid hydroperoxides in biological samples (*FRBM* Vol. 3, No. 5, 1987), which included contributions from seven research groups describing their methods of choice. Since that time we have received several more articles on the same topic, which we present here.

It may prove to be unusual to present two Forums on the same subject, but the detection of the oxidation products of lipids is an extremely important topic. Because radicals generally are very reactive and consequently short-lived, their direct detection is almost always impossible. Thus, indirect methods for detecting radical reactions are required. Even though it is true that electron spin resonance (ESR) in principle can detect radicals directly, the radicals dealt with in biological samples almost always are so reactive and short-lived that the indirect spin trap method must be used. Since the animal being hunted is so fleeting, one must look for its "footprints."

The "footprints" that free radical reactions leave in biological tissue are of a variety of types,¹ but it is remarkable how many involve searching for products of the oxidation of polyunsaturated fatty acids (PUFA). This is partly because of a historical bias, since PUFA oxidation was one of the first consequences of free radical reactions observed in biology, dating to the first observations of the rancidity of milk products and the thiobarbituric acid (TBA) test. Thus, the methods useful for detecting radical "footprints" include:

- ESR (directly, often in frozen samples)
- ESR spin trapping²⁻⁴
- TBA reactive materials (note these can form from PUFA and also from nucleic acids)⁵⁻⁹
- detection of malonaldehyde (MDA) by direct methods (such as HPLC of MDA itself or as its dinitrophenylhydrazone)^{10,11}
- detection of other oxidation products from PUFA (such as 4-hydroxynonenal)¹²
- measurement of lipid hydroperoxides¹³⁻¹⁹
- detection of volatile hydrocarbons (ethane, pentane, ethylene)²⁰⁻²²

- detection of oxidation products from lipids other than PUFA (e.g., cholesterol)^{23,24}
- oxidation of methional, methionine, or 2-keto-4-thiomethylbutanoic acid to ethylene^{25,26}
- oxidation of benzoic acid to carbon dioxide (often with radiolabelled carbon dioxide)²⁷
- oxidation of phenol, benzoic acid, or aspirin to hydroxylated products²⁸
- determination of decreases in antioxidant levels (e.g., decreased GSH, tocopherol, or ascorbate) or of increases in the oxidized products from antioxidants (e.g., tocopherol quinone or the ascorbyl radical)²⁹
- detection of oxidized DNA bases (e.g., thymine glycol, 8-hydroxydeoxyguanosine)³⁰
- detection of oxidized products from proteins (e.g., methionine sulfoxide from methionine) or of proteins oxidized to carbonyl-containing products that then react with hydride-reducing agents^{31,32}
- detection of adducts of DNA bases (e.g., by enzymatic hydrolysis post-labeling using ³²P)^{33,34}
- chemiluminescence methods^{35,36}

This is an impressive list indeed, but somehow one is left with a feeling that the "perfect, universal" method is not yet in hand. Maybe there is no perfect method, applicable for all situations. For example, smokers do not show elevated levels of TBA reactive substances (TBARS) in their plasma^{1,29} but do show elevated ethane/pentane, suggesting that one must search for oxidative changes in the organ at risk.²² Similarly, smokers have normal plasma vitamin E, but decreased levels in lung lavage.²⁹

It also is unfortunate that a single group does not more often publish the results of several of these tests, since, as indicated above, even a lack of agreement can be revealing. And agreement using different tests can serve to establish the validity and reliability of different indexes of oxidative stress.

Perhaps one of the greatest needs in the field now is the availability of a noninvasive test to probe the oxidative stress status (OSS) of humans. If we could measure OSS, we could ask the question: In what dis-

eases is OSS elevated? And then we could ask the \$64 questions: Do antioxidants prevent this elevation. And, if so, does this result in some protection against pathologies or diseases?

It is our hope that this Forum will focus attention of our readers on these and similar questions, and that you will find these articles useful and challenging. As always, your comments and suggestions are most welcome.

REFERENCES

1. Pryor, W. A. Vitamin E: The current research status and suggestions for future studies. *N.Y. Acad. Sci.* in press; 1989.
2. Janzen, E. G. A critical review of spin trapping in biological systems. In: Pryor, W. A., ed. *Free radicals in biology*. Vol. 4. New York: Academic Press; 1980:116-150.
3. McCay, P. B.; Noguchi, T.; Fong, K. L.; Lai, E. K.; Poyer, J. L. Production of radicals from enzyme systems and the use of spin traps. In: Pryor, W. A., ed. *Free radicals in biology*. Vol. 4. New York: Academic Press; 1980:155-184.
4. Mason, R. P. Free-radical intermediates in the metabolism of toxic chemicals. In: Pryor, W. A., ed. *Free radicals in biology*. Vol. 5. New York: Academic Press; 1980:161-212.
5. Kostrucha, J.; Kappus, H. Inverse relationship of ethane or n-pentane and malondialdehyde formed during lipid peroxidation in rat liver microsomes with different oxygen concentrations. *Biochem. Biophys. Acta* **879**:120-125; 1986.
6. Kosugi, H.; Kato, T.; Kikugawa, K. Formation of red pigment by a two-step 2-thiobarbituric acid reaction of alka-2,4-dienals. Potential products of lipid oxidation. *Lipids* **23**:1024; 1988.
7. Kosugi, H.; Kikugawa, K. Potential thiobarbituric acid-reactive substances in peroxidized lipids. *Free Radical Biol. Med.* **7**:205-207; 1989.
8. Yagi, K. A biochemical approach to atherogenesis. *Trends in Biochem. Sciences* **11**:18-19; 1986.
9. Gutteridge, J. M. C.; Rowley, D. A.; Halliwell, B. Superoxide-dependent formation of hydroxyl radicals in the presence of iron salts. *Biochem. J.* **199**:263-265; 1981.
10. Ekstrom, T.; Warholm, M.; Kronevi, T.; Hogberg, J. Recovery of malondialdehyde in urine as a 2,4-dinitrophenylhydrazine derivative after exposure to chloroform or hydroquinone. *Chem.-Biol. Interactions* **67**:25; 1988.
11. Dennis, K. J.; Shibamoto, T. Gas chromatographic determination of malonaldehyde formed by lipid peroxidation. *Free Radical Biol. Med.* **7**:187-192; 1989.
12. Esterbauer, H.; Zollner, H. Methods for determination of aldehydic lipid peroxidation products. *Free Radical Biol. Med.* **7**:900-900; 1989.
13. Smith, C. V.; Anderson, R. E. Methods for determination of lipid peroxidation in biological samples. *Free Radical Biol. Med.* **3**:341-344; 1987.
14. Terao, J.; Matsushita, S. Application of HPLC for the determination of phospholipid hydroperoxides in food and biological systems. *Free Radical Biol. Med.* **3**:345-348; 1987.
15. Van Kuijk, F. J. G. M.; Dratz, E. A. Detection of phospholipid peroxides in biological samples. *Free Radical Biol. Med.* **3**:349-354; 1987.
16. Pendleton, R. B.; Lands, W. E. M. Assay of lipid hydroperoxides by activation of cyclooxygenase activity. *Free Radical Biol. Med.* **3**:337-340; 1987.
17. Funk, M. O. The electroanalytical approach to lipid peroxide determinations. *Free Radical Biol. Med.* **3**:319-322; 1987.
18. Corongiu, F. P.; Banni, S.; Dessi, M. A. Conjugated dienes detected in tissue lipid extracts by second derivative spectrophotometry. *Free Radical Biol. Med.* **7**:183-186; 1989.
19. Piretti, M. V.; Pagliuca, G. Systematic isolation and identification of membrane lipid oxidation products. *Free Radical Biol. Med.* **7**:219-222; 1989.
20. Wendel, A. Measurement of in vivo lipid peroxidation and toxicological significance. *Free Radical Biol. Med.* **3**:355-358; 1987.
21. Dillard, C. J.; Tappel, A. L. Lipid peroxidation products in biological tissues. *Free Radical Biol. Med.* **7**:193-196; 1989.
22. Shariff, R.; Hoshino, E.; Allard, J.; Pichard, C.; Kurian, R.; Jeejeebhoy, K. N. Vitamin E supplementation in smokers. *Amer. J. Clin. Nutr.* **47**(4):758 abstr. (1988).
23. Smith, L. L.; Kulig, M. J.; Miller, D.; Ansari, G. A. S. Oxidation of cholesterol by dioxygen species. *J. Amer. Chem. Soc.* **100**:6206; 1978.
24. Addis, P. B.; Emanuel, H. A.; Bergmann, S. D.; Zavoral, J. H. Capillary GC quantification of cholesterol oxidation products in plasma lipoproteins of fasted humans. *Free Radical Biol. Med.* **7**:179-182; 1989.
25. Winston, G. W.; Cederbaum, A. I. NADPH-dependent production of oxy radicals by purified components of the rat liver mixed function oxidase system. *J. Biol. Chem.* **258**:1508-1513; 1983.
26. Pryor, W. A.; Tang, R. H. Ethylene formation from methional. *Biochem. Biophys. Res. Commun.* **81**:498-503; 1978.
27. Winston, G. W.; Cederbaum, A. I. Oxidative decarboxylation of benzoate to carbon dioxide by rat liver microsomes: a probe for oxygen radical production during microsomal electron transfer. *Biochemistry* **21**:4265; 1982.
28. Grootveld, M.; Halliwell, B. An aromatic hydroxylation assay for hydroxyl radicals utilizing high-performance liquid chromatography (HPLC). Use to investigate the effect of EDTA on the Fenton reaction. *Free Rad. Res. Comm.* **1**:243-250; 1986.
29. Pacht, E. R.; Kaseki, H.; Mohammed, J. R.; Cornwell, D. G.; Davis, B. Deficiency of vitamin E in the alveolar fluid of cigarette smokers. *J. Clin. Invest.* **77**:789-796; 1986.
30. Ames, B. N. Endogenous DNA damage as related to cancer and aging. *Mutation Res.* in press; 1989.
31. Brot, N.; Weissbach, H. Biochemistry and physiological role of methionine sulfoxide residues in proteins. *Arch. Biochem. Biophys.* **223**:271-281; 1983.
32. Stadtman, E. R. Oxidation of proteins by mixed-function oxidation systems: implications in protein turnover, ageing and neutrophil function. *Trends in Biochem. Sciences* **11**:11-12; 1986.
33. Phillips, D. H.; Hewer, A.; Martin, C. N.; Garner, R. C.; King, M. M. Correlation of DNA adduct levels in human lung with cigarette smoking. *Nature* **336**:790; 1988.
34. Randerath, E.; Mittal, D.; Randerath, K. Tissue distribution of covalent DNA damage in mice treated dermally with cigarette 'tar': preference for lung and heart DNA. *Carcinogenesis* **9**:75-80; 1988.
35. Iwaoka, T.; Tabata, F.; Takahashi, T. Lipid peroxidation and lipid peroxide detected by chemiluminescence. *Free Radical Biol. Med.* **3**:329-334; 1987.
36. Miyazawa, T. Determination of phospholipid hydroperoxides in human blood plasma by a chemiluminescence-HPLC assay. *Free Radical Biol. Med.* **7**:209-217; 1989.